

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problems Mailbox.**

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 7/00, 7/04, 15/00, A61K 39/21, 39/205	A1	(11) International Publication Number: WO 99/02657 (43) International Publication Date: 21 January 1999 (21.01.99)
(21) International Application Number: PCT/US98/14527 (22) International Filing Date: 10 July 1998 (10.07.98) (30) Priority Data: 60/052,366 11 July 1997 (11.07.97) US (71) Applicant (for all designated States except US): YALE UNIVERSITY [US/US]; Office of Cooperative Research, 155 Whitney Avenue, New Haven, CT 06520-8336 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): ROSE, John, K. [US/US]; 21 Shell Beach Road, Guilford, CT 06437 (US). SCHELL, Mathias [DE/US]; 1322 Town Walk Drive, Hamden, CT 06518 (US). JOHNSON, Erik, E. [US/US]; Apartment 3, 89 Linden Street, New Haven, CT 06511 (US). (74) Agent: KRINSKY, Mary, M.; 88 Prospect Street, New Haven, CT 06511-3797 (US).		(81) Designated States: AU, CA, JP, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: RHABDOVIRUSES WITH REENGINEERED COATS (57) Abstract Replication-competent recombinant rhabdoviruses that lack a functional glycoprotein gene and express at least one foreign polypeptide such as a cellular receptor for another virus in their viral envelopes are useful in the treatment of pathogenic viruses. In one embodiment, a recombinant vesicular stomatitis virus (VSV) lacking its glycoprotein (G) gene and expressing instead the HIV receptor and a coreceptor is employed in a method for treating persons infected with HIV. The recombinant virus is defective for entry into normal cells but is able to control HIV infection in a T cell line by replicating in, and killing, HIV-infected cells.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

RHABDOVIRUSES WITH REENGINEERED COATS

Related Application Data

This application claims priority benefit of co-pending U.S. application serial number 60/052,366, filed 11 July 1997.

Technical Field of the Invention

5 This invention relates primarily to replication-competent rhabdoviruses which lack a functional glycoprotein gene and express in the viral coat at least one receptor for another virus. Recombinant rhabdoviruses of the invention such as vesicular stomatitis virus which express at least one cellular receptor and a coreceptor of another pathogenic virus such as HIV are useful for the treatment of
10 patients infected with the pathogenic virus.

Background of the Invention

To cause infection, membrane-enveloped viruses such as human immunodeficiency virus type 1 (HIV-1) must first bind receptors at the cell surface and subsequently fuse their own membrane with that of the cell. This fusion process
15 releases the viral genetic material into the cytoplasm and initiates infection. Entry of HIV-1 into cells has long been known to require virus binding to the cell surface molecule CD4 (Maddon, *et al.*, 1986; Sattentau and Weiss, 1988), and recent studies have defined chemokine receptor molecules, CXCR4 (Feng, *et al.*, 1996; Berson, *et al.*, 1996), CCR5 (Alkhatib, *et al.*, 1996; Choe, *et al.*, 1996;

Deng, *et al.*, 1996; Doranz, *et al.*, 1996; Dragic, *et al.*, 1996), and CCR3 (Choe, *et al.*, 1996; Doranz, *et al.*, 1996) as co-receptors required for entry. HIV entry occurs in stages: initial binding of the viral envelope protein (designated gp120/41) to CD4 is followed by conformational changes that allow binding to
5 coreceptor and subsequent coalescence of the viral and cellular membranes (Lapham, *et al.*, 1996; Trkola, *et al.*, 1996; Wu, *et al.*, 1996). HIV-infected cells express gp120/41 on the cell surface during infection and prior to its incorporation into budding HIV-1 particles.

Recent studies have shown that vesicular stomatitis virus (VSV) has
10 potential as a high level expression vector capable of incorporating foreign proteins into the viral envelope (Schnell, *et al.*, 1996a; Schnell, *et al.*, 1996b). VSV causes an extremely rapid cytopathic infection in most animal cells including human T cells in culture, but is normally non-pathogenic in humans (for review see Wagner and Rose, 1996). Within two to three hours after infection, VSV
15 blocks host cell protein synthesis and within eight hours produces very large quantities of progeny virus particles that bud from the cell surface prior to cell lysis. VSV has a non-segmented, negative-strand RNA genome that is copied in the cytoplasm of infected cells by the viral RNA polymerase to generate five mRNAs encoding the five structural proteins. Only one of these proteins, a
20 glycoprotein designated G, is present in the viral membrane and is responsible for the very broad host range of VSV. G protein recognizes cell surfaces and catalyzes fusion of the viral membrane with cellular membranes (Florkiewicz and Rose, 1984). Foreign membrane proteins such as CD4 and other viral membrane proteins can be expressed at very high levels from the genome of recombinant
25 VSVs and these molecules are then incorporated at high levels into the viral membrane along with G protein (Schnell, *et al.*, 1996b).

Summary of the Invention

It is an objective of the invention to utilize this strategy to engineer recombinant viruses that can be employed to target cells infected with another virus such as HIV.

- 5 It is a further and more specific objective to provide a method for treating a person infected with HIV.

These and other objectives are accomplished by the present invention, which provides replication-competent recombinant rhabdoviruses such as vesicular stomatitis viruses (VSV) that lack a functional glycoprotein gene and express in
10 their viral coats at least one foreign polypeptide. In preferred embodiments, the foreign polypeptide is a cellular receptor for another virus and is expressed in a manner sufficient to target cells infected with the other virus. In one embodiment, the invention provides replication-competent recombinant vesicular stomatitis virus which lacks a functional G glycoprotein gene and displays at least one cellular
15 receptor for a pathogenic mammalian virus. Where the other virus is HIV, the expressed polypeptide is an HIV receptor such as CD4, an HIV coreceptor such as CXCR4, and/or mixtures of these.

The invention correspondingly provides methods for treating patients infected with a pathogenic virus by administering to the patient an effective
20 amount of a recombinant rhabdovirus of the invention such as VSV which expresses in its viral envelope at least one receptor for the pathogenic virus. As set out hereafter, treatment methods for patients infected with HIV-1 and/or HIV-2 are disclosed.

Description of the Figures

Figure 1 illustrates diagrams of recombinant VSV genomes. The order of the genes in wild-type VSV and in the recombinant VSVs described is shown. Each extra gene was inserted so that it contained the appropriate signals specifying VSV polymerase transcription initiation and termination. The details of plasmid constructions and derivation of the recombinant viruses are given in the Examples. The genes are transcribed from left to right from the negative strand RNA genome which is shown in the 3'-5' orientation.

Figure 2 shows photographs of cultures illustrating that VSVs lacking the G gene form plaques on a complementing cell line. BHK-G cells were seeded on 6-well plates (1.5×10^5 cells in each well), and G protein expression was induced by removal of tetracycline. The indicated dilutions of $100 \mu\text{l}$ the VSV Δ G-CD4 virus (10^{-2} , 10^{-3} and 10^{-4}) were added 16 h later, followed by addition of 2 ml of DME containing 2% methylcellulose. Cells were incubated for 48 h and then stained with naphthalene black to visualize the small plaques. Wild-type VSV plaques are much larger and are visible after 18h, but plaque formation by the defective virus on this cell line was slower, probably because of the reduced level of complementing VSV G protein expressed.

Figure 3 shows immunoelectron microscopic visualization of CD4 in the envelopes of VSV Δ G-CD4 and VSV Δ G-CC4. Purified wild-type VSV, VSV Δ G-CD4, or VSV Δ G-CC4 particles were negatively stained with phosphotungstic acid after labeling with a monoclonal antibody directed to CD4 followed by gold-conjugate goat anti-mouse IgG particles. Wild-type VSV particles contained visible spikes of VSV G protein trimers (arrows) and did not bind antibody to CD4 (wild-type VSV, left panel), while VSV Δ G-CD4 or VSV Δ G-CC4 bound anti-CD4 antibody and did not show visible spikes on the surface.

Figure 4 illustrates analysis of HIV-1 and VSVΔG-CC4 infection by immunofluorescence microscopy. Duplicate cultures of Jurkat cells were infected with HIV-1 strain IIIB and five days later, superinfected with VSVΔG-CC4 (panel D, E, F) or propagated without superinfection (panel A, B, C). Jurkat cells that
5 were uninfected with HIV-1 were also infected with VSVΔG-CC4 (panel G, H, I). Fourteen days after the VSVΔG-CC4 infection, cells were fixed and labeled with a mouse monoclonal antibody to VSV N protein and a secondary, affinity-purified, rhodamine-conjugated anti-mouse antibody (detected in panels C, F, I) and with a human polyclonal anti-HIV immune globulin followed by affinity-
10 purified, FITC-conjugated anti-human antibodies (detected in panels B, E, H). Panels A, D, and G show Nomarski images of the same fields of cells shown stained in the two images to their right.

Figure 5 provides line graphs illustrating quantitation of HIV-1 and VSVΔG-CC4 infection in Jurkat cells. Duplicate cultures of Jurkat cells were
15 infected with HIV-1 IIIB at day 0 and either superinfected with VSVΔ-CC4 five days later (■) or propagated without superinfection (□). The percentage of HIV⁺ cells was determined by immunofluorescence microscopy on the days indicated (Figure 5A). The percentage of HIV⁺ cells that were also infected with VSVΔG-CC4 was determined by immunofluorescence microscopy to detect VSV
20 N protein (Figure 5B). One hundred to two hundred cells were counted from multiple random fields to determine the percentages of cells staining positive for HIV or VSV antigens.

Figure 6 provides line graphs illustrating the effect of VSVΔG-CC4 infection on production of infectious HIV-1 and on release of RT. Media from
25 Jurkat cell cultures were analyzed for infectious HIV titers and for RT activity as described in the Examples. Titers (Figure 5A) were performed in triplicate and are reported as TCID₅₀/ml (□, HIV infected only; ■, HIV + VSVΔG-CC4). Titers determined by this assay were consistent with titers determined by an indirect immunofluorescence assay detecting HIV antigens. RT activity (Figure

5B) is reported in total counts per minute (cpm) bound to the DEAE paper after subtraction of a background of less than 50 cpm (□, HIV infected only; ■, HIV + VSVΔG-CC4; Δ, VSVΔG-CC4 infected only). The numbers plotted are the average of duplicates that agreed within 5%. Samples were collected from

5 Jurkat cultures on the indicated days after superinfection with VSVΔG-CC4.

Detailed Description of the Invention

This invention is based upon the construction of a new virus from vesicular stomatitis virus (VSV) that is replication-competent and kills cells rapidly, but is only able to infect cells that are first infected with HIV. The virus

10 contains a deletion in the VSV envelope glycoprotein gene and expresses instead the genes for the HIV receptor CD4 and an HIV coreceptor, CXCR4. The virus kills HIV-infected cells in culture and controls HIV infection.

In the practice of this invention, replication-competent recombinant rhabdoviruses are generated which lack a functional glycoprotein gene and express

15 a foreign polypeptide in their viral envelopes. By "foreign" is meant not naturally occurring in the rhabdovirus transformed. The foreign polypeptide is one that binds to a mammalian cell such as a cellular receptor and/or coreceptor for another virus in many embodiments, and an antibody in other embodiments. The invention is particularly suited to target cells infected with membrane-enveloped

20 viruses. The invention provides a general strategy of virus targeting to infected cells in cases where receptor(s) recognized by viral envelope proteins are known, and the viral protein has membrane fusion activity. Likewise, both conventional and hybrid antibodies to pathologic cells are also known and are used to target these cells. In either case, foreign polypeptides are expressed in the rhabdovirus

25 in a manner sufficient to target cells to be killed. Therefore, as used herein, by a "recombinant rhabdovirus lacking a functional glycoprotein gene" is meant a virus

having any alteration or disruption of the glycoprotein gene, and/or expressing a poorly functional or nonfunctional glycoprotein, or combinations thereof.

In the examples that follow, recombinant VSVs are generated which lack a functional envelope glycoprotein G or corresponding gene and express instead at least one receptor or coreceptor of an HIV virus. In preferred embodiments, the G gene is deleted, but any mutation of the gene that alters the host range specificity of VSV or otherwise eliminates the function of the G protein can be employed.

A gene for a foreign polypeptide such as one that binds to mammalian cells infected with a pathogenic virus is then inserted and/or added to the genome of the recombinant rhabdovirus. As summarized above, in many typical embodiments, the added gene is a receptor recognized by a virus glycoprotein and that is present on the surface of mammalian cells. However, recombinant rhabdoviruses engineered to kill other cells, *e.g.*, cancer cells, are encompassed by the invention. In these embodiments, genes for specific antibodies and a membrane fusion protein are typically incorporated into rhabdoviruses lacking the a functional glycoprotein gene.

The invention thus provides medical or veterinary methods for treating a patient (human being or animal) infected with a pathogenic mammalian virus by administration to the patient of an effective amount of a replication-competent recombinant vesicular stomatitis virus which lacks a functional G glycoprotein gene and displays a receptor of the pathogenic virus in its coat in a manner sufficient to target cells infected with the pathogenic virus. An example wherein the pathogenic, membrane-enveloped virus is HIV and the envelope proteins added to the VSV are an HIV receptor and a specific coreceptor is given in the Examples section hereafter. In that illustration, a T cell tropic HIV is employed, but recombinant VSVs of the invention can be engineered to express coreceptors of other HIV strains. Other embodiments express a receptor and not a coreceptor,

and some recombinant VSVs of the invention express more than one HIV coreceptor.

In the practice of a preferred embodiment of the invention, a patient
5 infected with HIV is treated by administering to the patient an effective amount of
a replication-competent recombinant vesicular stomatitis virus which lacks a
functional glycoprotein G gene and expresses at least one HIV receptor such as
CD4 and at least one HIV coreceptor such as CXCR4, CCR5, CCR3, and/or
mixtures thereof. An exemplary construct is VSVΔG-CC4 described hereafter and
10 illustrated in Figure 1.

Administration of a recombinant rhabdovirus of the invention to a person
or other mammal can be via any local or systemic method known by skilled
workers, but is preferably systemic. Systemic administration includes intravenous,
intramuscular, or intradermal administration by sterile injections, parenteral
15 administration, and the like, typically in combination with a pharmaceutically
acceptable carrier and/or other adjuvant or adjunct compound that maintains
viability of the recombinant VSV virus or other rhabdovirus, or enhances its
effect. Combinations of therapies may also be employed.

The amount of recombinant rhabdovirus necessary to bring about thera-
20 peutic treatment is not fixed *per se*, and necessarily depends upon the severity or
the extent of disease. Administration is facilitated and, in some cases additional
therapeutic effects are provided by, the carrier. When a carrier is employed, it is
necessary that it be inert in the sense of not inactivating the recombinant VSV or
other rhabdovirus and in the sense of not bringing about any adverse effect to the
25 patient to which it is administered. In most situations, doses are dependent upon
the extent of the patient's disease, the age, weight, and clinical condition of the
patient to be treated, the potency of the recombinant virus, the adjuvants or
adjunct compounds (if any) employed, and the concentrations of recombinant
rhabdovirus and other ingredients which are typically put together in association

with a pharmaceutically acceptable carrier. Most preferably, administration of the recombinant rhabdoviruses of the invention reduces the diseased cells in the patient so significantly that they are barely detectable or undetectable.

It is an advantage of the invention that preferred recombinant VSVs
5 which incorporate the receptor of one virus in the coat of another kill cells infected with the other virus rapidly and thus control infection. It is a further advantage of the invention that recombinant VSVs or other rhabdoviruses of the invention are genetically engineered to specific targets, and kill only infected or pathogenic cells.

10 The following examples are presented to further illustrate and explain the present invention and should not be taken as limiting in any regard.

Examples

In the studies reported herein VSV recombinants are generated which exhibit a complete deletion of the VSV G gene and a substitution instead of the
15 genes for CD4 and CXCR4. These viruses assemble particles containing both CD4 and CXCR4 in their envelopes and are targeted specifically to infect HIV-1-infected cells.

Experimental Procedures

Plasmid construction. Digestion with MluI and SphI was used to delete
20 the entire VSV G gene from pVSV-CD4 (Schnell et al., 1996b). The resulting plasmid was designated pVSVΔG-CD4. A plasmid encoding the CXCR4 protein with an HA epitope tag was described previously (Berson et al., 1996). CXCR4 was amplified by PCR from this plasmid with VENT polymerase (Stratagene) using primers 5'-ACTGCCCGGGCTCGAGGTTACCATGGAGGGGATCAG-3'

(SEQ ID NO: 1) and 5'-AGCTGCGGCCGCTAGCTTAGCTCCCGGGAAGAGATGCG-3' (SEQ ID NO: 2). The first primer contains an *Xho*I site and the second an *Nhe*I site (boldface letters). The PCR product was digested with *Xho*I and *Nhe*I and cloned into pVSV-XN1 (Schnell et al., 1996a) that was digested with *Xho*I and *Nhe*I. The resulting plasmid was designated pVSV-CXCR4. The G gene was deleted from pVSV-CXCR4 with *Xho*I and *Mlu*I. The coding sequence for human CD4 was amplified from pCD4 (Shaw et al., 1989) by PCR using primers 5'-CCGGGTACCA**CGCGTACAATGAACCGGGGAGTCCCTTTT**AG-3' (SEQ ID NO: 3) and 5'-GGGCCCCT**CGAGCGTGATATCTGT**TAGTTTTTTCATAC**TC**-AAATGGGGCTACATGTCTTC-3' (SEQ ID NO: 4). The first primer contains an *Mlu*I site (boldface letters); the second primer contains an *Xho*I site (boldface letters) and a VSV transcription stop-start signal (underlined letters). The CD4 gene was inserted at the site of the G deletion to generate a plasmid that was designated pVSVΔDG-CC4.

Construction of a BHK cell line expressing VSV G protein. The tetracycline-regulated system employed was described previously (Shockett et al., 1995). A VSV G gene was prepared from a plasmid designated pBSG by digestion with *Xho*I. Extensions were filled in with Klenow enzyme, and the fragment ligated to pTet-Splice (Life Technologies) that had been cleaved with *Eco*RV. The resulting plasmid was called pTET-G. BHK cells on 10 cm plates (20% confluent) were transfected with 1 μg pTET-G and 10 μg pTet-TAk (Life Technologies) using the CaPO₄ method (mammalian transfection kit; Stratagene). One day after transfection 0.5 μg/ml tetracycline was added to suppress G protein expression, which is toxic to BHK cells, and stable transfected cells were selected by adding 0.75 mg/ml geneticin (G418). Ten days after transfection, cell colonies were isolated and, after induction by removal of tetracycline, they were screened for expression of VSV G protein by immunofluorescence microscopy.

Recovery of VSV recombinants. Baby hamster kidney cells (BHK-21, ATCC) were maintained in DMEM (Dulbecco's modified Eagle's medium) supple-

mented with 5% fetal bovine serum (FBS). Cells on 10 cm dishes (~80% confluent) were infected at an MOI of 10 with vTF7-3 (Lawson et al., 1995). After 1 hour, plasmids encoding the N, P, G, and L proteins and the appropriate recombinant anti-genome RNA were transfected into the cells using a cationic liposome reagent containing dimethyldioctadecyl ammonium bromide and phosphatidylethanolamine (Lawson et al., 1995; Rose et al., 1991). Plasmid amounts were 10 μ g of the respective full length plasmid (VSV Δ G-CD4 or VSV Δ G-CC4), 3 μ g pBS-N, 5 μ g pBS-P, 4 μ g BS-G and 2 μ g pBS-L. Supernatants of recoveries were filtered through 0.22 μ m filters, and 5 ml was used to infect BHK-G cells on 10 cm plates which had been induced for twelve hours by removal of tetracycline. Supernatants of the first passage were filtered twice through a syringe filter (0.1 μ m pore size) to remove any vaccinia virus and the viruses were then passaged on 4 x 10⁶ BHK-G cells. Supernatants (10 ml) were titered on BHK-G cells and frozen at -80°C. For production of virus stocks, 2 x 10⁶ BHK cells on 10 cm plates were infected for 2 h with 10 ml supernatants of VSV Δ G-CD4 or VSV Δ G-CC4 grown on BHK-G cells, washed three times with phosphate buffered saline (PBS) followed by addition of 10 ml RPMI medium with 10% fetal bovine serum (FBS) or 10 ml DMEM with 5% FBS. At 16 h after infection, supernatants were clarified by centrifugation and used for experiments or stored at -80°C.

Preparation and analysis of viruses. For isolation of virions, a monolayer of BHK cells (~80% confluent) on a 10 cm dish was infected with VSV Δ G-CD4 or VSV Δ G-CC4 (multiplicity of infection, MOI=5), or wild-type VSV (MOI=0.01). At 16 h after infection, cell debris and nuclei were removed by centrifugation at 1,250 x g for 5 min and virus was then purified twice by ultracentrifugation at 120,000g through a 20% sucrose solution containing 10 mM Tris-HCl, pH 7.5, and 1 mM EDTA. Virus pellets were resuspended and analyzed by SDS-PAGE (10% acrylamide). Gels were either stained with Coomassie brilliant blue or transferred to nitrocellulose membranes. Membranes were blocked with 5% low fat milk powder in TBST (10 mM Tris-HCl pH 8.0, 150 mM NaCl, 0.05% Tween™-20) for 1 h at room temperature and then incubated with an anti-

body directed against CD4 (sheep α -CD4, AIDS Research Reference and Reagent Program) or an antibody 12CA5 (Boehringer-Mannheim) directed against the HA epitope in CXCR4. Proteins were visualized after incubation with horseradish peroxidase-conjugated donkey anti-sheep IgG or horseradish-peroxidase conjugated
5 goat anti-mouse IgG (Jackson Reasearch) for 1 h at room temperature using enhanced chemiluminescence (ECL; Amersham) system.

Electron microscopy and labeling with gold-conjugated antibodies.

Virus particles from infection of approximately 10^7 cells on a 15 cm dish were recovered from the culture medium by first pelleting cell debris at 1,500 x g for
10 10 min. The virus was then concentrated and purified by centrifugation as described above. Virus samples were absorbed onto carbon coated grids for 5 min and then blocked with 1% BSA in PBS for 10 min at room temperature. The grids were then placed on a 50 μ l drop of anti-CD4 Mab OKT4 (Reinherz et al., 1979) diluted 1:50 in PBS containing 1% BSA. After 1 hour, excess antibody was
15 removed by placing grids sequentially onto five 50- μ l drops of 1% BSA in PBS for 2 min each time. The grids were then placed on a 50- μ l drop of goat anti-mouse IgG (Fc) labeled with 15 nm gold particles (AuroProbe, Amersham Inc.). Unbound gold conjugates were removed by five sequential 2-min washes with PBS. The virus-immunogold complexes were then negative stained by incubation
20 the grids for 4 min on 50- μ l drops of 2% phosphotungstic acid (ph 7.2). Excess stain was removed and the grids were air dried. Images of viruses were obtained with a Zeiss EM910 electron microscope.

Infection and immunofluorescence microscopy of infected Jurkat cells.

Approximately 3×10^6 Jurkat Cells (clone E6-1, ATCC #TIB-125) were seeded in
25 T-25 flasks and grown in RPMI medium supplemented with 10% FBS. Infection with HIV-1 strain IIIB was at an MOI of approximately 0.01. At five days post-infection, cells were superinfected with 5 ml of medium containing VSV Δ G-CC4. As controls, one HIV-infected culture was not superinfected, and a third culture was infected only with VSV Δ G-CC4. Cells were collected for analysis at each

time point as follows. One half of the volume of the culture was collected and an equal volume of RPMI supplemented with 10% FBS was added back to the original cultures. Cells were pelleted at 1000 rpm for five minutes at 5°C in an IEC Centra-7R refrigerated centrifuge (Fisher Scientific). The media was
5 discarded, cells were washed once in 5 ml PBS, and pelleted again. The PBS was removed and cells were resuspended in 100 μ l PBS. The cells were then fixed in 1 ml of 3% paraformaldehyde for one hour at room temperature, or overnight at 4°C. After permeabilization in 1% Triton X-100, cells were stained with a
10 monoclonal antibody to VSV N protein and with human polyclonal HIV immunoglobulin (AIDS Research Reference and Reagent Program), followed by rhodamine-conjugated anti-mouse and FITC-conjugated anti-human antibodies (Jackson Research).

Cells were examined and photographed with 40X planapochromat objective on a Nikon Microphot-FX microscope equipped for epifluorescence.
15 Infection of cells with HIV-1 or VSV Δ G-CC4 was monitored by counting at least three random fields of cells that typically contained 100-200 cells. Syncytia were counted as single cells.

Determination of HIV Titers. HIV titers were obtained using supernatants from infected cultures and making serial three-fold dilutions in 96-well
20 plates. Ten thousand MT-2 cells provided by the AIDS Research Reference and Reagent Program (ARRRP) were added to each well and the virus and cells were mixed by pipeting. Plates were incubated at 37°C for 3-5 days and scored for the presence of syncytia. Titers at each time point were performed in triplicate and two or more syncytia per well were counted as positive for HIV infection.
25 Titers were calculated as described previously (Reed and Muench, 1938) and are reported as 50% tissue culture infectious doses (TCID₅₀) per ml.

Reverse Transcriptase Assays. Each assay was performed in 96-well plates in duplicate (Willey et al., 1988). Twenty-five microliters of reverse

transcription reaction mix (0.05% NP-40, 50 mM Tris-HCl, pH 7.8, 75 mM KCl, 2 mM DTT, 5 mM MgCl₂, 5 units oligo (dT)₁₂₋₁₈, 5 mg/ml poly A) was added, followed by addition of 10 μ l of cell-free culture supernatants. An additional 25 μ l of reaction mix was added containing 10 μ Ci/ml α -[³²P]-dTTP, and the reaction
5 was placed at 37°C for 90 minutes. Five microliters of each sample was then spotted onto Whatman DE81 paper, which was allowed to air dry. The DE81 paper was then washed in 2 x SSC (0.3 M NaCl, 20 mM sodium citrate) buffer four times, each time at room temperature for five minutes, followed by two one-minute washes in 95% ethanol at room temperature. After air drying, the paper
10 was exposed to a Molecular Dynamics PhosphorImager screen overnight. Samples were quantitated directly on the phosphorimager and by counting in a scintillation spectrometer after excision from the paper. Duplicate samples agreed within 5%.

Neutralization of infection by VSVDG-CC4. Polyclonal sheep anti-CD4 serum and anti-gp120 serum (ARRRP) were heat treated at 56°C for one hour to
15 inactivate complement. The anti-VSV neutralizing monoclonal antibody (I1) was a mouse ascites fluid. Neutralizations were performed as follows. VSV Δ G-CC4 virus (1 ml derived from $\sim 10^6$ infected BHK cells) was mixed with 100 μ l anti-CD4 or 100 μ l anti-VSV for one hour at 37°C, and then the mixture was added to $\sim 10^6$ Jurkat cells that had been infected with an MOI of 0.01 HIV-1 IIB five
20 days previously. Alternatively, 100 μ l anti-gp120 serum was added directly to 1 ml of HIV-infected Jurkat cells for one hour at 37°C. Then 1 ml of VSV Δ G-CC4 was added to the cells. After 3 days, cells were pelleted at 1000 rpm in a clinical centrifuge for 5 minutes and washed once in PBS. Cells were fixed and stained to detect HIV and VSV N protein antigens.

25

Results

Recovery of VSV recombinants expressing CD4 and CXCR4. Genetic engineering of non-segmented negative-strand RNA viruses such as VSV is complicated by the fact that the minimal infectious unit of the virus is not RNA,

but rather the viral RNA genome in a ribonucleoprotein complex composed of nucleocapsid (N) and polymerase proteins (L and P). Infectious VSV can be recovered from plasmid DNA as follows. Four separate plasmids expressing the full-length antigenomic VSV RNA and the N, P and L genes under control of bacteriophage T7 promoters are transfected into cells already infected with a vaccinia virus recombinant expressing T7 RNA polymerase (vTF7-3). Intracellular assembly of the antigenomic RNA into nucleocapsids is followed by VSV replication and transcription to generate infectious, replication-competent VSV (Fuerst et al., 1986; Lawson et al., 1995; Whelan et al., 1995). Additional genes can be expressed from extra transcription units introduced into the VSV genome (Schnell et al., 1996a).

The construction of a VSV recombinant expressing the CD4 protein in addition to the five VSV proteins N, P, M, G and L has been previously reported (Schnell et al., 1996b). To generate a recombinant expressing CD4 and CXCR4 instead of VSV G, the upstream VSV G gene was deleted from the VSVCD4 construct (Figure 1). The recovery of virus, designated VSVΔG-CD4, was then accomplished by including a complementing plasmid encoding VSV G protein in addition to those encoding N, P, and L. The gene encoding CXCR4 at the position downstream of CD4 was next added and the virus designated VSVΔG-CC4 was recovered. Because the recovered viruses are capable of only a single round of infection on BHK cells, they were initially recovered and grown on cells infected with a vaccinia virus encoding the T7 RNA polymerase and transiently expressing a complementing VSV G protein from a transfected plasmid DNA.

A cell line expressing VSV G protein allows growth of the defective viruses. In order to grow the VSVΔG-CD4 and VSVΔG-CC4 virus in the absence of vaccinia virus, a BHK cell line expressing the VSV G protein from an inducible promoter (Shockett et al., 1995) was generated. Figure 2 shows that the VSVΔG-CC4 virus forms plaques on this cell line indicating multiple rounds of infection. Titers obtained after growing the defective viruses on this line were in the range of

0.5 - 1×10^6 pfu/ml. These titers are reduced at least 1000-fold compared to those of wild-type VSV presumably because the levels of G protein provided by the complementing cell line are low compared to those expressed in wild-type infection.

5 Incorporation of CD4 and CXCR4 into the recombinant viruses. To determine if the CD4 and CXCR4 molecules were expressed from the defective viruses and incorporated into virus particles, BHK cells (not expressing VSV G) were infected with VSVΔG-CD4 or VSVΔG-CC4 that had been grown on the BHK-G line (MOI = 5). After 16 hours when all cells had been killed, the virus
10 particles were purified from the medium. Viral proteins were then separated by SDS-PAGE and detected by staining with Coomassie blue. The wild-type virus contains the L, G, N, and M proteins (P is a minor protein comigrating with N) while the VSVΔG-CD4 and VSVΔG-CC4 viruses lack G but contain a new protein of about 55,000 daltons, the size expected for CD4. The presence of CD4 in the
15 particles was also verified by western blotting with sheep anti-CD4 antibody or monoclonal antibody 12CA5 to the HA epitope tag in CXCR4. The CXCR4 protein was not detected in VSVΔG-CC4 by staining, perhaps because it contains seven membrane spanning segments that does not bind the stain, but was easily detected by western blotting.

20 Budding of VSV in the absence of VSV G protein is known to be approximately 20-fold less efficient than budding in its presence (Knipe et al., 1977). Similar results have been obtained for rabies virus (Mebatsion et al., 1996). Although the amounts of protein detected by Coomassie blue staining appeared similar for all viruses, the virus loaded in test lanes were derived from 10^7
25 infected BHK cells, while the wild-type virus in the control lane was from only 2×10^5 , or 50-fold fewer infected cells to achieve similar staining. The results not only confirm the importance of VSV G protein in driving highly efficient virus budding, but also illustrate that significant amounts of virus production occur even in the absence of VSV G protein.

To examine the morphology of the recombinant virus particles and to verify that CD4 was exposed on the viral membrane, purified virus particles were examined by electron microscopy after labeling with antibody to CD4 and a gold-conjugated secondary antibody. Photographs of negatively stained virus particles
5 are shown in Figure 3. Wild-type VSV particles contained visible G protein spikes (arrows) and did not bind antibody to CD4, while the VSVΔG-CD4 or VSVΔG-CC4 viruses lacked visible spikes and did bind anti-CD4 antibodies. The antibody to CXCR4 used in western blotting recognizes the internal C-terminal HA tag on CXCR4 and therefore is not suitable for labeling virus particles.

10 Infection of HIV-1 Infected Cells by VSVΔG-CC4. Because cells infected with HIV-1 have the gp 120/41 envelope protein on their surface, it might be possible to infect these cells with VSVΔG-CC4. The gp 120/41 molecules present on the cell surface should bind to CD4 and CXCR4 on the virus surface and promote fusion of the viral and cell membranes. To test this possibility, a
15 human T cell line (Jurkat) was infected with HIV-1 IIIB. At five days post-infection, it was expected from previous experiments that about half of the cells would be infected with HIV-1, and VSVΔG-CC4 virus was added. Two control cultures, one infected with HIV-1 only, and another infected with VSVΔG-CC4 only, were followed in parallel.

20 Initially infection by HIV-1 and VSVΔG-CC4 using indirect immunofluorescence microscopy to detect both HIV-1 proteins and VSV nucleocapsid (N) protein was followed. An example of results from the immunofluorescence analysis at 19 days after HIV infection is shown in Figure 4, along with a Nomar-ski image of the same field of cells. A compilation of the immunofluorescence
25 data obtained throughout the experiment is graphed in Figure 5. The fields shown in Figure 4 illustrate that in the control culture infected with HIV-1 only, nearly all cells stained positive for HIV antigens (A and B) while none of the cells stained positive for VSV N protein (C). In the cells infected with HIV-1 and VSVΔG-CC4, only a small fraction of the cells were positive for HIV-1 antigens (D and

E), and the majority of those cells were also positive for VSV N protein (F), indicating that they had been superinfected with VSVΔG-CC4. It was also noted that rare cells that were strongly positive for VSV N protein expression, had only little or no HIV protein evident. These cells may result from shut-off of cellular
5 and HIV protein synthesis by VSVΔG-CC4 or from other mechanisms of infection (see Discussion below). In Jurkat cells infected with VSVΔG-CC4 only, none of the cells showed the presence HIV or VSV N antigens (H, I, and J).

From the quantitation in Figure 5 it is evident that when cells were infected with HIV-1 alone, the percentage of HIV-1⁺ cells rose to nearly 100%
10 and remained high throughout the experiment. When cells were first infected with HIV-1 and subsequently infected with VSVΔG-CC4, the percentage of HIV⁺ cells began to decrease at day 14 and plateaued at day 21 to levels between 5 and 10% HIV⁺. In addition, Figure 5 shows that the percentage of HIV-1₊ cells that were also infected with VSVΔG-CC4 increased and then reached a plateau at around 90-
15 95%. It therefore appears that cells infected with HIV-1 are rapidly superinfected with VSVΔG-CC4 and killed before high levels of HIV-1 can be released. A balance is then reached between the two viruses where HIV infection is controlled by VSVΔG-CC4 infection and maintained at low levels. Very similar results were obtained in an experiment performed with the H9 T cell line, another cell line in
20 which HIV-1 IIIB grows well.

The possibility that the VSVΔG-CD4 virus lacking the coreceptor gene might infect HIV-1 infected cells was also tested. In one experiment it was observed that a low level of infection by VSVΔG-CD4 (about 10% that observed with VSVΔG-CC4) occurred. However, the initial infection could be completely
25 neutralized with antibody to VSV G, indicating that a trace of G protein sufficient to start infection must have been present in the virions. This G protein was presumably carried into the BHK cell line from the input virus grown in the BHK-G line and then incorporated into exiting virions. Interestingly, once the infection was established in HIV-infected Jurkat cells, it persisted for

several weeks, indicating that the virus could propagate without encoding its own coreceptor. Presumably budding virions can pick up cell-encoded coreceptor and are then able to propagate. However, control of HIV-1 infection was not as strong as with VSVΔG-CC4, indicating that it is preferable to have the virus
5 encode high levels of the coreceptor. Because the control of HIV infection was less effective without coreceptor, further studies with this virus were not pursued.

VSVΔG-CC4 infection controls release of infectious HIV-1. To obtain measurements of the effects VSVΔG-CC4 on production of infectious HIV-1 and on release of HIV-1 particles, the experiments shown in Figures 4 and 5 were
10 repeated. but instead followed HIV-1 titers and release of reverse transcriptase (RT) into the medium (Figures 6A and 6B). At the time of addition of VSVΔG-CC4 at three days after HIV-1 infection, HIV titers and RT levels in the medium were low. In the control cells infected with HIV-1 alone or in the cells super-
15 infected with VSVΔG-CC4, both the HIV titers and RT activity peaked at ten days after infection. There was then a decline in RT activity by day 14 in cells superinfected with VSVΔG-CC4 and a leveling off at about 10% of the RT level seen in control cells. The results with titers of infectious HIV-1 were much more
20 dramatic, with a complete disappearance of infectious HIV-1 (> 4 logs below the control) at day 14 followed by a fluctuation at levels of only 0-39 infectious particles per milliliter out to day 34. These titers should be compared to persistent titers of $\sim 10^4$ /ml for the control culture that was not infected with VSVΔG-CC4.

The RT results were consistent with what has been seen previously with the immunofluorescence in Figure 5, but the much larger decreases in infectious HIV titers were not expected. These lower titers could be explained if the
25 VSVΔG-CC4 infection selected for HIV-infected cells that expressed little or no HIV envelope and thus produced HIV particles lacking envelope but containing RT.

Anti-CD4 or anti-gp120 block infection by VSVΔG-CC4. Control neutralization experiments (Table 1) to verify that entry of VSVΔG-CC4 into HIV infected cells was dependent on CD4 and gp120 were also performed. VSVΔG-CC4 was incubated with antibody to CD4, neutralizing antibody to VSV-G, or without antibody before addition to Jurkat cells that had been infected with HIV-1 for five days as in Figure 5. Table 1 shows that anti-CD4 completely eliminated the VSVΔG-CC4 infection, while antibody to G did not.

Table 1. Neutralization of VSVΔG-CC4 Infectivity		
Antibody ^a	% HIV ⁺ Cells ^b	% VSV N ⁺ Cells ^b
None	60	39
anti-VSV G	59	37
anti-CD4	74	0
anti-HIV gp 120	71	0

^a Viruses grown on BHK cells not expressing VSV G were incubated with anti-VSV G or anti-CD4, or cells were incubated with anti-HIV gp120 prior to mixing of virus and cells.

^b HIV positive and VSV N⁺ cells were identified by indirect immunofluorescence microscopy as in Figures 4 and 5. Approximately 100 cells from random fields were counted for each determination. The percentage of cells positive for HIV antigens or VSV N protein is indicated.

Similarly, pre-incubation of the HIV-infected cells with a polyclonal anti-gp120 antibody completely prevented infection by VSVΔG-CC4. Because CD4 can bind MHC class II on cell surfaces (Doyle and Strominger, 1987), it was conceivable that some low level of infection could occur after binding of VSVΔG-CC4 to MHC class II on cells, although this would not be expected without a membrane fusion protein. To test this possibility Daudi cells, a lymphocyte cell line known to express abundant quantities of MHC class II protein (Till et al., 1988), was

used. These cells were readily infected by wild-type VSV, but no infection by VSVΔG-CC4 was observed.

Discussion

A recombinant VSV that lacks its own envelope protein gene (G) and
5 expresses instead the HIV-1 receptor and coreceptor molecules, CD4 and CXCR4
was constructed. This novel virus is defective for entry into normal cells, but can
be propagated on cells expressing a complementing VSV envelope protein. The
complemented virus can then be grown on cells not expressing G protein where it
buds particles containing CD4 and CXCR4 in the viral membrane. These viruses
10 are replication competent, but can enter and propagate only in cells that are first
infected with HIV-1 and express HIV-1 envelope protein. Infection of these cells
by VSVΔG-CC4 is neutralized by antibody to CD4 added to the virus, or by
antibody to HIV gp120 added to the cells.

The results showed that in a mixed population of cells about half of
15 which were HIV-1-infected and half were uninfected, superinfection with VSVΔG-
CC4 ultimately limited production of infectious HIV-1 to levels that were barely
detectable or undetectable, at least 300-fold to 104-fold lower than HIV titers
produced by control cells that were not superinfected with VSVΔG-CC4. This
reduction presumably is the result of killing of the HIV-1 infected cells by the
20 VSVΔG-CC4 virus. The reduction in HIV-1 titers was greater than would have
been predicted from analysis of the number of HIV-1-infected cells or from RT
assays. The number of HIV positive cells was reduced about ten-fold by the
VSVΔG-CC4 infection, and the release of RT was reduced to a similar extent. In
considering the mechanism of VSVΔG-CC4 action, this discrepancy is reasonable
25 since VSVΔG-CC4 will target and kill those cells expressing HIV envelope protein
and thus select for cells that produce little or no HIV envelope. Such cells would
still score positive for HIV-1 antigen production, and would produce normal levels

of HIV-like particles containing RT, but the majority of these particles would be noninfectious.

The persistence of a low level of both HIV-1 and VSVΔG-CC4 coinfection for over one month in culture indicates that both viruses are continuing to replicate and that they have reached a balance. Based on the data, it seems unlikely that VSVΔG-CC4 will ever completely eliminate the HIV-1 infection in the culture. This low level persistence may occur because VSVΔG-CC4 can only infect HIV-1 infected cells after gp120/41 is expressed on the cell surface and such cells likely release some HIV-1 before they are killed.

10

The VSVΔG-CC4 virus infects cells that have already been infected with HIV-1, but it may also be fusing directly with HIV-1 virions. Such fused viruses would contain the nucleocapsids of HIV and VSV and might be able to initiate infection if sufficient gp120/41 were still present on the membrane. Such an infection would almost certainly favor replication of VSVΔG-CC4 because VSV would shut off cellular protein synthesis in 2-3 hours, even before HIV had sufficient time for reverse transcription and integration. Infection by these fused viruses could explain the presence of cells (Figure 4) that are making VSV proteins but do not appear to have synthesized HIV proteins. While not wishing to be bound to any theory, it is also possible that VSV has inhibited HIV protein synthesis in these cells. Infection of uninfected cells by VSVΔG-CC4 might also occur if VSVΔG-CC4 pseudotypes carrying HIV envelope protein were generated in HIV-infected cells. Incorporation of the wild-type HIV envelope into VSV particles is extremely inefficient (Johnson et al., 1997) and therefore this latter mechanism would likely be rare. To test directly for such pseudotypes, a short-term infection of Jurkat cells with supernatants from cells coinfecting with HIV-1 and VSVΔG-CC4 was carried out. VSV protein expression was not detected in these cells, while it was readily detected by using the same supernatant to infect HIV-infected Jurkat cells. It was concluded that the VSVΔG-CC4(HIV) pseudotypes are extremely rare if they exist at all.

30

Several aspects of this system that are worth noting. First, because VSVΔG-CC4 is specifically targeted to cells already infected with HIV-1, the VSVΔG-CC4 infection level would be expected to parallel HIV infection and disappear with clearance of HIV. Second, VSVΔG-CC4 encodes only human proteins as its envelope proteins, and thus would not induce production of neutralizing antibodies. The internal VSV proteins would be expected to induce cytotoxic T cell responses, but such responses might help in clearing HIV infection by rapidly killing cells infected with HIV-1 and VSVΔG-CC4. Third, mutations to drug resistance are rapidly selected in HIV-1, but resistance to infection by VSVΔG-CC4 would require loss of HIV's ability to bind CD4 or coreceptor and would therefore not be selected.

The results also illustrate an important aspect of VSV budding and assembly. Based on virus protein yields and titers, it is estimated that each BHK cell infected with wild-type VSV produces approximately 100,000 particles, and up to 10% of these particles are infectious. The deletion of the VSV G gene reduced particle yields of VSVΔG-CC4 or VSVΔG-CD4 approximately 30-50 fold, indicating a substantial role for G in promoting budding. However, despite this reduced budding, at least 2,000 particles of VSVΔG-CC4 or VSVΔG-CD4 are produced from each BHK cell in the absence of G protein. The incorporation of CD4 and CXCR4 into these particles is presumed to occur by a passive process of protein trapping as virus buds from the cell surface, because neither protein would be expected to contain specific signals promoting incorporation. Because VSV blocks host protein synthesis and expresses extremely large amounts of its mRNA and proteins, CD4 and CXCR4 are likely to be the major proteins present on cell surface and therefore the major proteins in the VSVΔG-CC4 envelope.

References

Alkhatib, G., Combadiere, C., Broder, C. C., Feng, Y., Kennedy, P. E., Murphy, P. M., and Berger, E. A. (1996). CC CKR5: a RANTES, MIP-1alpha,

- MIP-1beta receptor as a fusion cofactor for macrophage-tropic HIV-1. *Science* 272, 1955-8.
- 5 Berson, J. F., Long, D., Doranz, B. J., Rucker, J., Jirik, F. R., and Doms, R. W. (1996). A seven-transmembrane domain receptor involved in fusion and entry of T-cell-tropic human immunodeficiency virus type 1 strains. *J Virol* 70, 6288-95.
- 10 Choe, H., Farzan, M., Sun, Y., Sullivan, N., Rollins, B., Ponath, P. D., Wu, L., Mackay, C. R., LaRosa, G., Newman, W., Gerard, N., Gerard, C., and Sodroski, J. (1996). The beta-chemokine receptors CCR3 and CCR5 facilitate infection by primary HIV-1 isolates. *Cell* 85, 1135-48.
- Deng, H., Liu, R., Ellmeier, W., Choe, S., Unutmaz, D., Burkhart, M., Di Marzio, P., Marmon, S., Sutton, R. E., Hill, C. M., Davis, C. B., Peiper, S. C., Schall, T. J., Littman, D. R., and Landau, N. R. (1996). Identification of a major co-receptor for primary isolates of HIV-1. *Nature* 381, 661-6.
- 15 Doranz, B. J., Rucker, J., Yi, Y., Smyth, R. J., Samson, M., Peiper, S. C., Parmentier, M., Collman, R. G., and Doms, R. W. (1996). A dual-tropic primary HIV-1 isolate that uses fusin and the beta-chemokine receptors CKR-5, CKR-3, and CKR-2b as fusion cofactors. *Cell* 85, 1149-58.
- 20 Doyle, C., and Strominger, J. L. (1987). Interaction between CD4 and class II MHC molecules mediates cell adhesion. *Nature* 330, 256-9.
- Dragic, T., Litwin, V., Allaway, G. P., Martin, S. R., Huang, Y., Nagashima, K. A., Cayanan, C., Maddon, P. J., Koup, R. A., Moore, J. P., and Paxton, W. A. (1996). HIV-1 entry into CD4+ cells is mediated by the chemokine receptor CC-CKR-5. *Nature* 381, 667-73.
- 25 Feng, Y., Broder, C. C., Kennedy, P. E., and Berger, E. A. (1996). HIV-1 entry cofactor: functional cDNA cloning of a seven-transmembrane, G protein-coupled receptor. *Science* 272, 872-7.
- Florkiewicz, R. Z., and Rose, J. K. (1984). A cell line expressing vesicular stomatitis virus glycoprotein fuses at low pH. *Science* 225, 721-3.
- 30 Fuerst, T. R., Niles, E. G., Studier, F. W., and Moss, B. (1986). Eukaryotic transient-expression system based on recombinant vaccinia virus that synthesizes bacteriophage T7 RNA polymerase. *Proc. Natl. Acad. Sci. USA* 83, 8122-6.
- Johnson, J. E., Schnell, M. J., Buonocore, L., and Rose, J. K. (1997). Specific targeting to CD4+ Cells of recombinant vesicular stomatitis viruses encoding human immunodeficiency virus envelope proteins. *J. Virol.* 71, 5060-5068.
- 35

- Knipe, D. M., Baltimore, D., and Lodish, H. F. (1977). Maturation of viral proteins in cells infected with temperature-sensitive mutants of vesicular stomatitis virus. *J. Virol.* 21, 1149-58.
- 5 Lapham, C. K., Ouyang, J., Chandrasekhar, B., Nguyen, N. Y., Dimitrov, D. S., and Golding, H. (1996). Evidence for cell-surface association between fusin and the CD4-gp120 complex in human cell lines. *Science* 274, 602-5.
- Lawson, N. D., Stillman, E. A., Whitt, M. A., and Rose, J. K. (1995). Recombinant vesicular stomatitis viruses from DNA. *Proc. Natl. Acad. Sci. U S A* 92, 4477-81.
- 10 Maddon, P. J., Dalgleish, A. G., McDougal, J. S., Clapham, P. R., Weiss, R. A., and Axel, R. (1986). The T4 gene encodes the AIDS virus receptor and is expressed in the immune system and the brain. *Cell* 47, 333-48.
- Mebatsion, T., Konig, M., and Conzelmann, K. K. (1996). Budding of rabies virus particles in the absence of the spike glycoprotein. *Cell* 84, 941-951.
- 15 Moore, J. P. (1997). Coreceptors: implications for HIV pathogenesis and therapy. *Science* 276, 51-52.
- Reed, L. J., and Muench, H. (1938). A Simple Method of Estimating Fifty Percent Endpoints. *American Journal of Hygiene* 27, 493-497.
- 20 Reinherz, E. L., Kung, P. C., Goldstein, G., and Schlossman, S. F. (1979). Separation of functional subsets of human T cells by a monoclonal antibody. *Proc. Natl. Acad. Sci. USA* 76, 4061-5.
- Riedel, H., Kondor-Koch, C., and Garoff, H. (1984). Cell surface expression of fusogenic vesicular stomatitis virus G protein from cloned cDNA. *Embo J* 3, 1477-83.
- 25 Rose, J. K., Buonocore, L., and Whitt, M. A. (1991). A new cationic liposome reagent mediating nearly quantitative transfection of animal cells. *Biotechniques* 10, 520-5.
- Sattentau, Q. J., and Weiss, R. A. (1988). The CD4 antigen: physiological ligand and HIV receptor. *Cell* 52, 631-3.
- 30 Schnell, M. J., Buonocore, L., Kretzschmar, E., Johnson, E., and Rose, J. K. (1996b). Foreign glycoproteins expressed from recombinant vesicular stomatitis viruses are incorporated efficiently into virus particles. *Proc. Natl. Acad. Sci. USA* 93, 11359-11365.

- Schnell, M. J., Buonocore, L., Whitt, M. A., and Rose, J. K. (1996a). The minimal conserved transcription stop-start signal promotes stable expression of a foreign gene in vesicular stomatitis virus. *J. Virol.* 70, 2318-2323.
- 5 Schuitemaker, H., and Miedema, F. (1996). Viral and cellular determinants of HIV-1 replication in macrophages. *Aids* 10, S25-32.
- Shaw, A. S., Amrein, K. E., Hammond, C., Stern, D. F., Sefton, B. M., and Rose, J. K. (1989). The lck tyrosine protein kinase interacts with the cytoplasmic tail of the CD4 glycoprotein through its unique amino-terminal domain. *Cell* 59, 627-36.
- 10 Shockett, P., Difilippantonio, M., Hellman, N., and Schatz, D. G. (1995). A modified tetracycline-regulated system provides autoregulatory, inducible gene expression in cultured cells and transgenic mice. *Proc Natl Acad Sci U S A* 92, 6522-6.
- 15 Till, M. A., Ghetie, V., Gregory, T., Patzer, E. J., Porter, J. P., Uhr, J. W., Capon, D. J., and Vitetta, E. S. (1988). HIV-infected cells are killed by rCD4-ricin A chain. *Science* 242, 1166-8.
- 20 Trkola, A., Dragic, T., Arthos, J., Binley, J. M., Olson, W. C., Allaway, G. P., Cheng-Mayer, C., Robinson, J., Maddon, P. J., and Moore, J. P. (1996). CD4-dependent, antibody-sensitive interactions between HIV-1 and its co-receptor CCR-5 [see comments]. *Nature* 384, 184-7.
- Wagner, R. R., and Rose, J. K. (1996). Rhabdoviridae: The viruses and their replication. In *Fields Virology*, B. N. Fields and D. M. Knipe, eds., (New York: Lippincott-Raven).
- Weiss, R. A. (1993). How does HIV cause AIDS? *Science* 260, 1273-9.
- 25 Whelan, S. P. J., Ball, L. A., Barr, J. N., and Wertz, G. T. W. (1995). Efficient recovery of infectious vesicular stomatitis virus entirely from cDNA clones. *Proc. Natl. Acad. Sci. USA* 92, 8388-8392.
- 30 Willey, R. L., Smith, D. H., Lasky, L. A., Theodore, T. S., Earl, P. L., Moss, B., Capon, D. J., and Martin, M. A. (1988). In vitro mutagenesis identifies a region within the envelope gene of the human immunodeficiency virus that is critical for infectivity. *J. Virol.* 62, 139-147.
- 35 Wu, L., Gerard, N. P., Wyatt, R., Choe, H., Parolin, C., Ruffing, N., Borsetti, A., Cardoso, A. A., Desjardin, E., Newman, W., Gerard, C., and Sodroski, J. (1996). CD4-induced interaction of primary HIV-1 gp120 glycoproteins with the chemokine receptor CCR-5. *Nature* 384, 179-83.

The papers cited herein are expressly incorporated in their entireties by reference.

The invention was made with partial government support under NIH grants AI24345 and AI49235. The government has certain rights in the invention.

- 5 The above description is for the purpose of teaching the person of ordinary skill in the art how to practice the present invention, and it is not intended to detail all those obvious modifications and variations of it which will become apparent to the skilled worker upon reading the description. It is intended, however, that all such obvious modifications and variations be included within the
- 10 scope of the present invention, which is defined by the following claims. The claims are intended to cover the claimed components and steps in any sequence which is effective to meet the objectives there intended, unless the context specifically indicates the contrary.



CLAIMS

1. A replication-competent recombinant rhabdovirus which lacks a functional glycoprotein gene and encodes at least one foreign polypeptide that is expressed in its viral envelope.
2. A recombinant virus according to claim 1 wherein the polypeptide is a cellular receptor of another virus which is expressed in the rhabdovirus in a manner sufficient to target cells infected with the other virus.
3. A recombinant virus according to claim 1 which is a vesicular stomatitis virus which lacks a functional G glycoprotein gene and encodes at least one cellular receptor of a pathogenic mammalian virus.
4. A recombinant virus according to claim 3 wherein the pathogenic virus is HIV-1 or HIV-2.
5. A recombinant virus according to claim 4 wherein the pathogenic virus is HIV-1.
6. A recombinant virus according to claim 4 which expresses in its viral envelope at least one HIV receptor, at least one HIV co-receptor, or mixtures thereof.
7. A recombinant virus according to claim 3 comprising the additional genomes illustrated in Figure 1.
8. A method for treating a patient infected with a pathogenic virus comprising administering to the patient an effective amount of a recombinant virus according to claims 1, 2, 3, 4, 5, 6, or 7 which expresses in its viral envelope at least one receptor of the pathogenic virus so that the patient's cells harboring the pathogenic virus are targeted.

9. A replication-competent recombinant vesicular stomatitis virus which lacks a functional glycoprotein G gene and expresses at least one HIV receptor and at least one HIV co-receptor.

10. A method for treating a patient infected with HIV-1 or HIV-2 by administering to the patient an effective amount of a recombinant virus according to claims 1, 2, 3, 4, 5, 6, 7, or 9.

1/4

**Figure 1****Figure 2**

2/4




Figure 3








Figure 4

**Figure 5A****Figure 5B**

**Figure 6A****Figure 6B**

1/3

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANTS: John K. Rose
Matthias Schnell
Erik Johnson
- (ii) TITLE OF INVENTION: Rhabdoviruses with Reengineered Coats
- (iii) NUMBER OF SEQUENCES: 4
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESS: Yale School of Medicine
 - (B) STREET: 310 Cedar Street Street
 - (C) CITY: New Haven
 - (D) STATE: Connecticut
 - (E) COUNTRY: United States of America
 - (F) ZIP CODE: 06510
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: 3.5" 1.44 Mb diskette
 - (B) COMPUTER: IBM PC
 - (C) OPERATING SYSTEM: MS DOS
 - (D) SOFTWARE: Word Processing
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER: PCT/US/98/
 - (B) FILING DATE: 10-JUL-1998
 - (D) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:
 - (A) APPLICATION NUMBER: 60/052,366
 - (B) FILING DATE: 11-JUL-1997
- (viii) ATTORNEY INFORMATION:
 - (A) NAME: Mary M. Krinsky
 - (B) REGISTRATION NO.: 32423
 - (C) REFERENCE/DOCKET NUMBER: OCR-882
- (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE NUMBER: 203-773-9544
 - (B) TELEFAX NUMBER: 203-772-0587

2/3

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 36
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE:

- (A) DESCRIPTION: DNA

(ix) FEATURE:

- (A) NAME/KEY: primer used in constructs

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

ACTGCCCGGG CTCGAGGTTA CCATGGAGGG GATCAG

36

(3) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 38
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE:

- (A) DESCRIPTION: DNA

(ix) FEATURE:

- (A) NAME/KEY: primer used in constructs

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

AGCTGCGGCC GCTAGCTTAG CTCCCGGGAA GAGATGCG

38

(4) INFORMATION FOR SEQ ID NO: 3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 41
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE:

- (A) DESCRIPTION: DNA

(ix) FEATURE:

- (A) NAME/KEY: primer used in constructs

3/3

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

CCGGGTACCA CGCGTACAAT GAACCGGGGA GTCCCTTTTA G 41

(5) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 61
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE:

- (A) DESCRIPTION: DNA

(ix) FEATURE:

- (A) NAME/KEY: primer used in constructs

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

GGGCCCCCTCG AGCGTGATAT CTGTTAGTTT TTTTCATACT 40

CAAATGGGGC TACATGTCTT C 61

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US98/14527

A. CLASSIFICATION OF SUBJECT MATTER IPC(6) : C12N 7/00, 7/04, 15/00; A61K 39/21, 39/205 US CL : 435/235.1, 236, 320.1; 424/199.1, 224.1 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) U.S. : 435/235.1, 236, 320.1; 424/199.1, 224.1 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) USPATFUL, WPIDS, MEDLINE		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	LAWSON, N.D. et al., Recombinant vesicular stomatitis viruses from DNA. Proc. Natl. Acad. Sci. USA. May 1995, Vol. 92, pages 4477-4481, especially page 4478.	1-7 and 9
Y	MEBATSION, T. et al., Highly stable expression of a foreign gene from rabies virus vectors. Proc. Natl. Acad. Sci. USA. July 1996, Vol. 93, pages 7310-7314, especially page 7311.	1-7 and 9
Y	SCHNELL, M. J. et al., Foreign glycoproteins expressed from recombinant vesicular stomatitis viruses are incorporated efficiently into virus particles. Proc. Natl. Acad. Sci. USA. October 1996, Vol. 93, pages 11359-11365, especially 11360.	1-7 and 9
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents:	"T"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be of particular relevance	"X"	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"B" earlier document published on or after the international filing date	"Y"	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Z"	document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means		
"P" document published prior to the international filing date but later than the priority date claimed		
Date of the actual completion of the international search	Date of mailing of the international search report	
28 AUGUST 1998	17 SEP 1998	
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231	Authorized officer <i>Jeffrey S. Parkin</i> Jeffrey S. Parkin, Ph.D.	
Facsimile No. (703) 305-3230	Telephone No. (703) 308-0196	

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/14527

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	FENG, Y. et al., HIV-1 entry cofactor: functional cDNA cloning of a seven-transmembrane G protein-coupled receptor. Science. 10 May 1996, Vol. 272, pages 872-877, see entire document.	6